



| Drug Classification | Street/Trade Names | Approx Window of Detection | Screen Cutoff ng/ml | Method of Use | Duration of Use | Possible Effects | Possible Effects of Overdose |
|------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------|---------------------|------------------------------------------------------|------------------|------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|
| Amphetamines: Includes Amphetamine and Methamphetamine (<i>Stimulant</i>) | Crystal, Ecstasy* (MDMA), Crank, Speed, Meth, Bennies, Uppers, Black beauties, Desoxyyn, Adderall, Dexedrine, MDA, "E", "X", Pane, | 1 - 2 Days (Longer for high use) | 1000 | Oral Injected Snorted Smoked (Freebased) | 2-4 Hours | Increased alertness, excitation, euphoria, increased pulse rate, loss of appetite, insomnia | Agitation, increased body temperature, hallucinations, convulsions, possible death |
| Barbiturates (<i>Depressant</i>) | Reds, Seconal, Downers, Goofballs, Amytal, Butisol, Fiorinal, Nembutal, Phenobarbital, Tuinal | 1 Day- short acting 2-3 Weeks - long acting | 200 | Oral, Injected | 1-16 Hours | Slurred speech, disorientation, Loss of inhibitions – memory, Impaired judgement | Shallow respiration, cold & clammy skin, weak & rapid pulse, coma, possible death |
| Benzodiazepines (<i>Depressant</i>) | Tranks, Ativan, Dalmane, Diazepam, Librium, Lorazepam, Valium, Clonazepam, Halcion, Prazepam, Oxazepam, Xanax | 3 - 5 Days (Up to 3 weeks, or longer for prolonged use) | 200 | Oral, Injected | 4-8 Hours | Slurred speech, disorientation, loss of inhibitions – memory, Impaired judgement | Shallow respiration, cold & clammy skin, weak & rapid pulse, coma, possible death |
| Cocaine (<i>Stimulant</i>) | Coke, Flake, Snow, Crack, Nose Candy | 2 - 4 Days (Up to 7 days or longer for chronic high use) | 300 | Snorted Injected Smoked (Freebased) | 1-2 Hours | Increased alertness, excitation, euphoria, increased pulse rate, loss of appetite, insomnia | Agitation, increased body temperature, hallucinations, convulsions, possible death |
| Ethanol [Alcohol] (<i>Depressant</i>) | Alcohol, Booze, Liquor, Spirits, Wine | 2 - 14 Hours | 20 mg/dl | Oral | 1 Hour per drink | Slurred speech, disorientation, loss of inhibitions – memory, impaired judgement | Shallow respiration, cold & clammy skin, weak & rapid pulse, coma, possible death |
| Ethylglucuronide (<i>Ethanol Metabolite</i>) | N/A – unique metabolite resulting from ethanol exposure | Up to 96 hours | 250 | Oral | N/A | See Ethanol | See Ethanol |
| Methadone (<i>Narcotic</i>) | Synthetic narcotic, Dolophine, Methadose, Dome | 3 Days | 300 | Oral Injected | 12-24 Hours | Euphoria, drowsiness, respiratory depression, constricted pupils, nausea | Slow & shallow breathing, clammy skin, convulsions, coma, possible death |
| Opiates [Codeine-Morphine-Heroin & synthetics] (<i>Narcotic</i>) | Smack, Horse, Big M, Dovers powder, Robitussin A-C, Tylenol w/codeine MS Contin, Dura-morph, Vicodin, Hydrocodone, Dilaudid Oxycontin* (Oxycodone) | 1 - 3 Days | 300 | Oral Injected Smoked | 3-6 Hours | Euphoria, drowsiness, respiratory depression, constricted pupils, nausea | Slow & shallow breathing, clammy skin, convulsions, coma, possible death |
| Phencyclidine [PCP] (<i>Hallucinogen</i>) | PCP, Angel Dust, Hog | 5 - 8 Days (Up to 30 days for chronic use) | 25 | Oral Injected Smoked | Variable | Illusions, poor perception of time & distance, hallucinations | Longer & more intense "trip" episodes, awake coma, bizarre behavior, psychosis, possible death |
| Propoxyphene (<i>Narcotic</i>) | Darvon, Darvocet | 1 - 2 Days | 300 | Oral | Variable | Euphoria, drowsiness, respiratory depression, constricted pupils, nausea | Slow & shallow breathing, clammy skin, convulsions, coma, possible death |
| THC (Marijuana) (<i>Cannabis</i>) | Weed, Pot, Grass, Gold, MJ, Bud | 5 days moderate use 10 Days heavy use Up to 2 months heavy chronic use | 20 or 50 | Oral Smoked | 2-4 Hours | Euphoria, lack of motivation, short term memory loss, lowered productivity, increased appetite | Fatigue, paranoia. |

Note: This information is intended to be used as a guideline and should not be applied to any particular individual. sales@regtox.com

© SRL Rev 7-2007

* **CONTACT STERLING** for information on our **Ecstasy & Oxycodone High Specificity Testing.**