

# Oral Fluid Drug Screen Chart

Compliments of  
(800) 442-0438



**STERLING**  
REFERENCE LABORATORIES  
a division of regional toxicology services, llc

Drug Classification	Street/Trade Names	Approximate Window of Detection <sup>1</sup>	Screen Cutoff ng/ml	Methods of Use	Duration of Use	Possible Effects	Possible Effects of Overdose
<b>Amphetamine</b> (Stimulant)	Speed, Bennies, Uppers, Adderall, Dexedrine, MDA, Vyvance	Up to 2 days	100	Oral, Injected, Snorted, Smoked (Freebased)	2-4 hours	Increased alertness, excitation, euphoria, increased pulse rate, loss of appetite, insomnia	Agitation, increased body temperature, hallucinations, convulsions
<b>Barbiturates</b> (Depressant)	Black beauties, Seconal, Reds, Downers, Goofballs, Amytal, Butisol, Fiorinal, Nembutal, Phenobarbital, Tuinal	Up to 2 days	20	Oral, injected	1-16 hours	Slurred speech, disorientation, loss of inhibitions – memory, impaired judgment	Shallow respiration, cold & clammy skin, weak & rapid pulse, coma, possible death
<b>Benzodiazepines</b> (Depressant)	Tranks, Ativan, Diazepam, Librium, Lorazepam, Valium, Clonazepam, Halcion, Prazepam, Oxazepam, Xanax	Up to 2 days	1.0	Oral, Injected	4-8 hours	Same as above	Same as above
<b>Cocaine</b> (Stimulant)	Coke, Flake, Snow, Crack	Up to 3 days	20	Snorted, Injected, Smoked (Freebased)	1-2 hours	Increased alertness, excitation, euphoria, increased pulse rate, loss of appetite, insomnia	Agitation, increased body temperature, hallucinations, convulsions
<b>Methadone</b> (Narcotic)	Synthetic narcotic, Dolophine, Methadose, Dome	Up to 2 days for occasional use Up to 3-5 days for chronic use	5	Oral, Injected	12-24 hours	Euphoria, drowsiness, respiratory depression, constricted pupils, nausea	Slow & shallow breathing, clammy skin, convulsions, coma, possible death
<b>Methamphetamine</b> (Stimulant)	Crystal, Crank, Ecstasy, Meth, MDMA, Uppers, Desoxyn	Up to 2 days	50	Oral, Injected, Snorted, Smoked (Freebased)	2-4 hours	Increased alertness, excitation, euphoria, increased pulse rate, loss of appetite, insomnia	Agitation, increased body temperature, hallucinations, convulsions
<b>Opiates</b> [Codeine-Morphine-Heroin] (Narcotic)	Smack, Horse, Big M, Dovers powder, Robitussin A-C, Tylenol w/codeine, Hydrocodone, MS Contin, Vicodin, Dura-morph	Up to 2 days	10	Oral, Injected, Smoked	3-6 hours	Euphoria, drowsiness, respiratory depression, constricted pupils, nausea	Slow & shallow breathing, clammy skin, convulsions, coma, possible death
<b>Oxycodone</b>	Percocet, Percodan, Oxycontin	Up to 2 days	20	Same as above	3-6 hours	Same as above	Same as above
<b>Phencyclidine (PCP)</b> (Hallucinogen)	PCP, Angel Dust, Hog	Up to 2 days	1.0	Oral, Injected, Smoked	variable	Illusions, poor perception of time & distance, hallucinations	Longer & more intense "trip" episodes, awake coma, bizarre behavior, psychosis, possible death
<b>Propoxyphene</b>	Darvon, Darvocet	Up to 1 day	20	Oral	variable	Euphoria, drowsiness, respiratory depression, constricted pupils, nausea	Slow & shallow breathing, clammy skin, convulsions, coma, possible death
<b>THC (Marijuana)</b> (Cannabis)	Weed, Pot, Grass, Gold	Up to 2 days	1.0	Oral, Smoked	2-4 hours	Euphoria, lack of motivation, short term memory loss, lowered productivity, increased appetite	Fatigue, paranoia

<sup>1</sup> Compilation of data is from the Journal of Analytical Toxicology publications, laboratory partners and other third correspondences. Data is specific to LABORATORY-BASED ORAL FLUID Screen. RAPID ORAL FLUID SCREENS TEND TO HAVE SHORTER WINDOW OF DETECTION, PARTICULARLY FOR MARIJUANA. \*\*Note: This information is intended to be used as a guideline and should not be applied to any particular individual. \*\* For certain drug classes that have many drugs included, window of detection may vary for different drugs.